

Map Unit Description

Weld County, Colorado, Northern Part

4 Ascalon fine sandy loam, 0 to 6 percent slopes

Setting

Elevation: 4500 to 6500 feet
Mean annual precipitation: 13 to 17 inches
Mean annual air temperature: 46 to 57 degrees F
Frost-free period: 130 to 160 days

Composition

Ascalon and similar soils: 85 percent
Minor components: 15 percent

Description of Ascalon

Setting

Landform: Plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous loamy alluvium

Properties and Qualities

Slope: 0 to 6 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.60 to 2.00 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 10 percent
Gypsum maximum: 0 percent
Available water capacity: Moderate (about 6.9 inches)

Interpretive Groups

Land capability classification (irrigated): 3e
Land capability (non irrigated): 3e
Ecological site: Loamy Plains (R067BY002CO)

Typical Profile

0 to 8 inches: fine sandy loam
8 to 22 inches: sandy clay loam
22 to 60 inches: sandy loam

Minor Components

Olney

Percent of map unit: 8 percent

Otero

Percent of map unit: 7 percent

Map Unit Description

Weld County, Colorado, Northern Part

36 Manzanola clay loam, 0 to 3 percent slopes

Setting

Elevation: 4400 to 5600 feet
Mean annual precipitation: 11 to 15 inches
Mean annual air temperature: 46 to 52 degrees F
Frost-free period: 140 to 180 days

Composition

Manzanola and similar soils: 85 percent
Minor components: 15 percent

Description of Manzanola

Setting

Landform: Stream terraces, swales, plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous clayey alluvium

Properties and Qualities

Slope: 0 to 3 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low or moderately high (0.06 to 0.20 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 5 percent
Gypsum maximum: 3 percent
Sodium adsorption ratio maximum: 15.0
Available water capacity: High (about 9.6 inches)

Interpretive Groups

Land capability (non irrigated): 4e
Ecological site: Clayey Plains (R067BY042CO)

Typical Profile

0 to 3 inches: clay loam
3 to 25 inches: clay
25 to 48 inches: clay
48 to 60 inches: clay loam

Minor Components

Avar

Percent of map unit: 15 percent

KUMMER PL LE23-99H2
T8N-R61W SEC 23: NWNW

Map Unit Description

Weld County, Colorado, Northern Part

45 Olney fine sandy loam, 6 to 9 percent slopes

Setting

Elevation: 3500 to 5800 feet
Mean annual precipitation: 11 to 15 inches
Mean annual air temperature: 46 to 54 degrees F
Frost-free period: 125 to 175 days

Composition

Olney and similar soils: 85 percent
Minor components: 15 percent

Description of Olney

Setting

Landform: Plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous loamy alluvium

Properties and Qualities

Slope: 6 to 9 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.57 to 2.00 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 0 percent
Available water capacity: Moderate (about 8.1 inches)

Interpretive Groups

Land capability (non irrigated): 6e
Ecological site: Loamy Plains (R067BY002CO)

Typical Profile

0 to 6 inches: fine sandy loam
6 to 18 inches: sandy clay loam
18 to 60 inches: sandy loam
60 to 64 inches: sandy loam

Minor Components

Stoneham

Percent of map unit: 5 percent

Vona

Percent of map unit: 5 percent

Ascalon

Percent of map unit: 5 percent