

Land USX Y31-01
Production Facility
T2N, R64W 6th P.M.
Section 31: NW NE

Map Unit Description

Weld County, Colorado, Southern Part

25 Haverson loam, 0 to 1 percent slopes

Setting

Elevation: 4500 to 4800 feet
Mean annual precipitation: 12 to 17 inches
Mean annual air temperature: 46 to 54 degrees F
Frost-free period: 125 to 180 days

Composition

Haverson and similar soils: 85 percent
Minor components: 15 percent

Description of Haverson

Setting

Landform: Flood plains, stream terraces
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Stratified, calcareous alluvium

Properties and Qualities

Slope: 0 to 1 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.60 to 2.00 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 1 percent
Available water capacity: High (about 9.6 inches)

Interpretive Groups

Land capability classification (irrigated): 3w
Ecological site: Loamy Plains (R067BY002CO)

Typical Profile

0 to 4 inches: loam
4 to 60 inches: stratified loamy sand to loam to clay loam

Minor Components

Vona

Percent of map unit: 8 percent

Fluvaquentic haplustolls

Percent of map unit: 4 percent
Landform: Terraces

Other soils

Percent of map unit: 3 percent

Land USX Y31-01
Production Facility
T2N, R64W 6th P.M.
Section 31: NWNE

Map Unit Description

Weld County, Colorado, Southern Part

18 Colby-Adena loams, 3 to 9 percent slopes

Setting

Elevation: 4750 to 4900 feet
Mean annual precipitation: 12 to 16 inches
Mean annual air temperature: 48 to 55 degrees F
Frost-free period: 120 to 160 days

Composition

Colby and similar soils: 55 percent
Adena and similar soils: 30 percent
Minor components: 15 percent

Description of Colby

Setting

Landform: Ridges, hills, plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous eolian deposits

Properties and Qualities

Slope: 5 to 9 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.57 to 2.00 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 0 percent
Available water capacity: High (about 10.6 inches)

Interpretive Groups

Land capability (non irrigated): 4e
Ecological site: Loamy Slopes (R067BY008CO)

Typical Profile

0 to 7 inches: loam
7 to 60 inches: silt loam

Description of Adena

Setting

Landform: Hills, plains, ridges
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous eolian deposits

Properties and Qualities

Slope: 3 to 7 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.60 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 0 percent
Available water capacity: High (about 10.1 inches)

Interpretive Groups

Land capability classification (irrigated): 3e
Land capability (non irrigated): 4e
Ecological site: Loamy Plains (R067BY002CO)

Typical Profile

0 to 6 inches: loam
6 to 9 inches: clay loam
9 to 60 inches: silt loam

Land USX Y31-01
Production Facility
T2N, R64W 6th P.M.
Section 31: NWNE

Map Unit Description

Weld County, Colorado, Southern Part

Minor Components

Kim
Percent of map unit: 5 percent

Keith
Percent of map unit: 4 percent

Wiley
Percent of map unit: 3 percent

Weld
Percent of map unit: 3 percent

Land USX Y31-01
Production Facility
T2N, R64W 6th P.M.
Section 31: NWNE

Map Unit Description

Weld County, Colorado, Southern Part

48 Olney fine sandy loam, 3 to 5 percent slopes

Setting

Elevation: 4600 to 5200 feet
Mean annual precipitation: 11 to 15 inches
Mean annual air temperature: 46 to 54 degrees F
Frost-free period: 125 to 175 days

Composition

Olney and similar soils: 85 percent
Minor components: 15 percent

Description of Olney

Setting

Landform: Plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Mixed deposit outwash

Properties and Qualities

Slope: 3 to 5 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.57 to 2.00 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 0 percent
Available water capacity: Moderate (about 7.0 inches)

Interpretive Groups

Land capability classification (irrigated): 3e
Land capability (non irrigated): 4c
Ecological site: Sandy Plains (R067BY024CO)

Typical Profile

0 to 10 inches: fine sandy loam
10 to 20 inches: sandy clay loam
20 to 25 inches: sandy clay loam
25 to 60 inches: fine sandy loam

Minor Components

Zigweid

Percent of map unit: 9 percent

Vona

Percent of map unit: 6 percent