

Map Unit Description

Weld County, Colorado, Northern Part

4 Ascalon fine sandy loam, 0 to 6 percent slopes

Setting

Elevation: 4500 to 6500 feet
Mean annual precipitation: 13 to 17 inches
Mean annual air temperature: 46 to 57 degrees F
Frost-free period: 130 to 160 days

Composition

Ascalon and similar soils: 85 percent
Minor components: 15 percent

Description of Ascalon

Setting

Landform: Plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous loamy alluvium

Properties and Qualities

Slope: 0 to 6 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.60 to 2.00 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 10 percent
Gypsum maximum: 0 percent
Available water capacity: Moderate (about 6.9 inches)

Interpretive Groups

Land capability classification (irrigated): 3e
Land capability (non irrigated): 3e
Ecological site: Loamy Plains (R067BY002CO)

Typical Profile

0 to 8 inches: fine sandy loam
8 to 22 inches: sandy clay loam
22 to 60 inches: sandy loam

Minor Components

Olney

Percent of map unit: 8 percent

Otero

Percent of map unit: 7 percent

Map Unit Description

Weld County, Colorado, Northern Part

5 Ascalon fine sandy loam, 6 to 9 percent slopes

Setting

Elevation: 4500 to 6500 feet
Mean annual precipitation: 13 to 17 inches
Mean annual air temperature: 46 to 57 degrees F
Frost-free period: 130 to 160 days

Composition

Ascalon and similar soils: 85 percent
Minor components: 15 percent

Description of Ascalon

Setting

Landform: Plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous loamy alluvium

Properties and Qualities

Slope: 6 to 9 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.60 to 2.00 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 10 percent
Gypsum maximum: 0 percent
Available water capacity: Moderate (about 6.8 inches)

Interpretive Groups

Land capability classification (irrigated): 4e
Land capability (non irrigated): 4e
Ecological site: Loamy Plains (R067BY002CO)

Typical Profile

0 to 6 inches: fine sandy loam
6 to 21 inches: sandy clay loam
21 to 60 inches: sandy loam

Minor Components

Altvan

Percent of map unit: 7 percent

Peetz

Percent of map unit: 4 percent

Cascajo

Percent of map unit: 3 percent

Aquic haplustolls

Percent of map unit: 1 percent
Landform: Swales

Map Unit Description

Weld County, Colorado, Northern Part

65 Terry sandy loam, 3 to 9 percent slopes

Setting

Elevation: 4000 to 6500 feet
Mean annual precipitation: 13 to 15 inches
Mean annual air temperature: 46 to 48 degrees F
Frost-free period: 120 to 180 days

Composition

Terry and similar soils: 85 percent
Minor components: 15 percent

Description of Terry

Setting

Landform: Plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous sandy residuum weathered from sandstone

Properties and Qualities

Slope: 3 to 9 percent
Depth to restrictive feature: 20 to 40 inches to Paralithic bedrock
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low or high (0.06 to 2.00 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 0 percent
Available water capacity: Low (about 3.7 inches)

Interpretive Groups

Land capability (non irrigated): 6e
Ecological site: Sandy Plains (R067BY024CO)

Typical Profile

0 to 5 inches: sandy loam
5 to 17 inches: fine sandy loam, sandy loam
17 to 32 inches: fine sandy loam, sandy loam, gravelly sandy loam
32 to 36 inches: weathered bedrock

Minor Components

Tassel

Percent of map unit: 5 percent

Olney

Percent of map unit: 4 percent

Renohill

Percent of map unit: 3 percent

Vona

Percent of map unit: 3 percent