

Lincoln County, Colorado

109—Ascalon-Haxtun complex, 0 to 3 percent slopes

Map Unit Setting

National map unit symbol: 3j8d
Elevation: 4,400 to 6,000 feet
Mean annual precipitation: 14 to 16 inches
Mean annual air temperature: 46 to 52 degrees F
Frost-free period: 135 to 155 days
Farmland classification: Not prime farmland

Map Unit Composition

Ascalon and similar soils: 55 percent
Haxtun and similar soils: 30 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Ascalon

Setting

Landform: Plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Eolian deposits

Typical profile

Ap - 0 to 4 inches: sandy loam
Bt - 4 to 15 inches: sandy clay loam
Bk - 15 to 60 inches: fine sandy loam

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat):
Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum in profile: 15 percent
Salinity, maximum in profile: Nonsaline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: Moderate (about 8.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3c
Hydrologic Soil Group: B
Ecological site: Sandy plains (R067BY024CO)

Description of Haxtun

Setting

Landform: Drainageways
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Alluvium and/or eolian deposits

Typical profile

Ap - 0 to 4 inches: loamy sand
BA - 4 to 17 inches: sandy loam
Bt - 17 to 44 inches: sandy clay loam
Btkb - 44 to 60 inches: silt loam

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat):
Moderately high (0.20 to 0.60 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum in profile: 15 percent
Gypsum, maximum in profile: 2 percent
Salinity, maximum in profile: Nonsaline (0.0 to 2.0 mmhos/cm)
Available water storage in profile: Moderate (about 8.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: B
Ecological site: Sandy plains (R067BY024CO)

Minor Components

Olne1

Percent of map unit: 3 percent
Landform: Plains
Other vegetative classification: SANDY PLAINS (067XY024CO_1)

Otero

Percent of map unit: 3 percent
Landform: Hills
Landform position (three-dimensional): Side slope
Other vegetative classification: SANDY PLAINS (067XY024CO_1)

Pleasant

Percent of map unit: 3 percent
Landform: Depressions
Ecological site: Plains swale (R067XY010CO)

Platner

Percent of map unit: 3 percent

Landform: Plains

Ecological site: Loamy plains (R067BY002CO)

Other vegetative classification: LOAMY PLAINS (067XY002CO_1)

Vona

Percent of map unit: 3 percent

Landform: Hills

Landform position (three-dimensional): Side slope

Other vegetative classification: SANDY PLAINS (067XY024CO_1)

Data Source Information

Soil Survey Area: Lincoln County, Colorado

Survey Area Data: Version 11, Dec 23, 2013

Lincoln County, Colorado

194—Valent-Bijou loamy sands, 1 to 12 percent slopes

Map Unit Setting

National map unit symbol: 2t50f
Elevation: 4,000 to 5,500 feet
Mean annual precipitation: 13 to 16 inches
Mean annual air temperature: 46 to 54 degrees F
Frost-free period: 125 to 170 days
Farmland classification: Not prime farmland

Map Unit Composition

Valent, warm, and similar soils: 50 percent
Bijou, warm, and similar soils: 40 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Valent, Warm

Setting

Landform: Ridges, hills
Landform position (two-dimensional): Backslope, summit
Landform position (three-dimensional): Side slope, head slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Eolian sands derived from sedimentary rock

Typical profile

A - 0 to 7 inches: loamy sand
AC - 7 to 16 inches: loamy sand
C - 16 to 79 inches: sand

Properties and qualities

Slope: 1 to 15 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Excessively drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.99 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline (0.1 to 0.2 mmhos/cm)
Sodium adsorption ratio, maximum in profile: 0.1
Available water storage in profile: Low (about 4.1 inches)

Interpretive groups

Land capability classification (irrigated): 6e
Land capability classification (nonirrigated): 6e
Hydrologic Soil Group: A
Ecological site: Deep sand (R067BY015CO)

Other vegetative classification: DEEP SANDS (067XY015CO_2)

Description of Bijou, Warm

Setting

Landform: Ridges, hills

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Linear

Across-slope shape: Convex, linear

Parent material: Eolian deposits derived from sedimentary rock

Typical profile

A - 0 to 4 inches: loamy sand

AB - 4 to 9 inches: loamy sand

Bt - 9 to 36 inches: sandy loam

BC - 36 to 50 inches: loamy sand

C - 50 to 79 inches: loamy sand

Properties and qualities

Slope: 1 to 9 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Somewhat excessively drained

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Salinity, maximum in profile: Nonsaline (0.0 to 0.1 mmhos/cm)

Available water storage in profile: Low (about 5.5 inches)

Interpretive groups

Land capability classification (irrigated): 4e

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: A

Ecological site: Sandy plains (R067BY024CO)

Minor Components

Vona, warm

Percent of map unit: 5 percent

Landform: Hills, ridges

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Linear

Across-slope shape: Convex, linear

Ecological site: Sandy plains (R067BY024CO)

Other vegetative classification: Unnamed (G067BW019CO)

Olnest

Percent of map unit: 5 percent

Landform: Hills, interfluvies

Landform position (two-dimensional): Summit, footslope

Landform position (three-dimensional): Side slope, interfluvie

Down-slope shape: Linear
Across-slope shape: Convex
Ecological site: Sandy plains (R067BY024CO)
Other vegetative classification: Unnamed (G067BW019CO)

Data Source Information

Soil Survey Area: Lincoln County, Colorado
Survey Area Data: Version 11, Dec 23, 2013